Table 5.4: Alcohol Consumption (Percent)^{1,2}

Alcohol Consumption	Total	Age Group			Disability Level		
						ADL Difficulty	
		65-74	75-84	85+	Moderate ³	Receives No Help	Receives Help
All Participants	(N=1002)	(N=388)	(N=311)	(N=303)	(N=343)	(N=478)	(N=181)
Do you usually drink alcoholic beverages, including beer, wine, sherry, or liquor, at least once every week? Yes	15.6	16.9	14.5	14.5	17.2	14.9	14.0
Current Alcohol Drinkers ⁴	(N=154)	(N=65)	(N=44)	(N=45)	(N=59)	(N=72)	(N=23)
Average drinks per week ⁵ 1-3 4-7 8 or more	36.3 37.5 26.2	34.4 38.2 27.4	36.7 36.8 26.4	41.6 37.3 21.1	38.0 40.5 21.5	32.5 33.8 33.6	43.0 41.2 15.9
On the days when you drink, about how many drinks do you usually have? 1 2 3 or more	54.6 30.1 15.3	47.3 29.8 23.0	60.9 29.7 9.4	62.0 32.2 5.8	56.8 26.0 17.2	47.1 35.8 17.1	71.0 23.5 5.6
Over the past 6 months, how many days per week did you typically drink like this? 1-2 3-4 5 or more	39.0 22.3 38.7	44.3 19.7 36.0	30.1 27.5 42.4	45.1 16.7 38.2	35.3 28.1 36.6	39.6 17.1 43.3	46.1 22.9 31.0

(Women's Health and Aging Study, baseline interview, 1992-1995)

All variables have less than 4% missing data. Results are based on non-missing data.

Descriptive statistics are based on weighted data.

No ADL difficulty; disabled in two or more domains (see Chapter 1).

Categories for each item may not add up to 100% due to rounding.

d(w), where d=drinks per day (On the days when you drink, about how many drinks do you usually have?) and w=days (Over the past 6 months, how many days per week did you typically drink like this?)